

## Product datasheet for RC215335L3V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Protein Kinase D2 (PRKD2) (NM 016457) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Protein Kinase D2 (PRKD2) (NM\_016457) Human Tagged ORF Clone Lentiviral Particle

**Symbol:** Protein Kinase D2

**Synonyms:** HSPC187; nPKC-D2; PKD2

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 016457

ORF Size: 2634 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC215335).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 016457.3

RefSeq Size: 2883 bp
RefSeq ORF: 2637 bp
Locus ID: 25865
UniProt ID: Q9BZL6
Cytogenetics: 19q13.32

**Protein Families:** Druggable Genome, Protein Kinase

**MW:** 96.5 kDa





# Protein Kinase D2 (PRKD2) (NM\_016457) Human Tagged ORF Clone Lentiviral Particle – RC215335L3V

#### **Gene Summary:**

The protein encoded by this gene belongs to the protein kinase D (PKD) family of serine/threonine protein kinases. This kinase can be activated by phorbol esters as well as by gastrin via the cholecystokinin B receptor (CCKBR) in gastric cancer cells. It can bind to diacylglycerol (DAG) in the trans-Golgi network (TGN) and may regulate basolateral membrane protein exit from TGN. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]