

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC215366L3V


## ACK1 (TNK2) (NM\_001010938) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	ACK1 (TNK2) (NM_001010938) Human Tagged ORF Clone Lentiviral Particle
Symbol:	TNK2
Synonyms:	ACK; ACK-1; ACK1; p21cdc42Hs
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001010938
ORF Size:	3258 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC215366).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001010938.1, NP 001010938.1</u>
RefSeq Size:	4222 bp
RefSeq ORF:	3144 bp
Locus ID:	10188
UniProt ID:	<u>Q07912</u>
Cytogenetics:	3q29
Protein Families:	Druggable Genome, Protein Kinase
MW:	119.35 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits<br/>both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs.<br/>This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3<br/>domain. The protein may be involved in a regulatory mechanism that sustains the GTP-<br/>bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation<br/>signal transduction pathway. Several alternatively spliced transcript variants have been<br/>identified from this gene, but the full-length nature of only two transcript variants has been<br/>determined. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US