

Product datasheet for RC215374L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GAB2 (NM_012296) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GAB2 (NM 012296) Human Tagged ORF Clone Lentiviral Particle

Symbol: GAB2

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_012296

ORF Size: 2028 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC215374).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 012296.2</u>

 RefSeq Size:
 6062 bp

 RefSeq ORF:
 1917 bp

 Locus ID:
 9846

 UniProt ID:
 Q9UQC2

Cytogenetics: 11q14.1

Protein Families: Druggable Genome

Protein Pathways: Chronic myeloid leukemia, Fc epsilon RI signaling pathway, Fc gamma R-mediated

phagocytosis





ORÏGENE

MW:

74.5 kDa

Gene Summary:

This gene is a member of the GRB2-associated binding protein (GAB) gene family. These proteins contain pleckstrin homology (PH) domain, and bind SHP2 tyrosine phosphatase and GRB2 adapter protein. They act as adapters for transmitting various signals in response to stimuli through cytokine and growth factor receptors, and T- and B-cell antigen receptors. The protein encoded by this gene is the principal activator of phosphatidylinositol-3 kinase in response to activation of the high affinity IgE receptor. Two alternatively spliced transcripts encoding different isoforms have been described for this gene. [provided by RefSeq, Nov 2009]