

### **Product datasheet for RC215420**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL22 RA2 (IL22RA2) (NM\_181310) Human Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: IL22 RA2 (IL22RA2) (NM\_181310) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: IL22RA2

Synonyms: CRF2-10; CRF2-S1; CRF2X; IL-22BP; IL-22R-alpha-2; IL-22RA2; ZCYTOR16

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RC215420 representing NM\_181310

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGATGCCTAAACATTGCTTTCTAGGCTTCCTCATCAGTTTCTTCCTTACTGGTGTAGCAGGAACTCAGT CAACGCATGAGTCTCTGAAGCCTCAGAGGGTACAATTTCAGTCCCGAAATTTTCACAACATTTTGCAATG GCAGCCTGGGAGGGCACTTACTGGCAACAGCAGTGTCTATTTTGTGCAGTACAAAATATATGGACAGAGA CAATGGAAAAATAAAGAAGACTGTTGGGGTACTCAAGAACTCTCTTGTGACCTTACCAGTGAAACCTCAG ACATACAGGAACCTTATTACGGGAGGGTGAGGGCGGCCTCGGCTGGGAGCTACTCAGAATGGAGCATGAC

GCCGCGGTTCACTCCCTGGTGGGAAAGAGCAAAAGGTTTA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC215420 representing NM\_181310

Red=Cloning site Green=Tags(s)

MMPKHCFLGFLISFFLTGVAGTQSTHESLKPQRVQFQSRNFHNILQWQPGRALTGNSSVYFVQYKIYGQR

QWKNKEDCWGTQELSCDLTSETSDIQEPYYGRVRAASAGSYSEWSMTPRFTPWWERAKGL

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 


**Restriction Sites:** Sgfl-Mlul






### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

### Plasmid Map:



**ACCN:** NM\_181310

ORF Size: 390 bp



**OTI Disclaimer:** 

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** 

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 181310.2</u>

 RefSeq Size:
 2644 bp

 RefSeq ORF:
 393 bp

 Locus ID:
 116379

 UniProt ID:
 Q969|5

 Cytogenetics:
 6q23.3

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

**MW:** 14.9 kDa

**Gene Summary:** This gene encodes a member of the class II cytokine receptor family. The encoded soluble

protein specifically binds to and inhibits interleukin 22 activity by blocking the interaction of interleukin 22 with its cell surface receptor. The encoded protein may be important in the regulation of inflammatory response, and has been implicated in the regulation of

tumorigenesis in the colon. Alternative splicing results in multiple transcript variants encoding

distinct isoforms. [provided by RefSeg, Jul 2013]