

Product datasheet for **RC215420**

IL22 RA2 (IL22RA2) (NM_181310) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	IL22 RA2 (IL22RA2) (NM_181310) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	IL22 RA2
Synonyms:	CRF2-10; CRF2-S1; CRF2X; IL-22BP; IL-22R-alpha-2; IL-22RA2; ZCYTOR16
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<p>>RC215420 representing NM_181310</p> <p>Red=Cloning site Blue=ORF Green=Tags(s)</p> <p>TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC</p> <p>ATGATGCCTAAACATTGCTTTCTAGGCTTCCTCATCAGTTTCTTCCTTACTGGTGTAGCAGGAAGTCACTCAACGCATGAGTCTCTGAAGCCTCAGAGGGTACAATTTTCAGTCCCGAAATTTTCACAACATTTTGCAATGGCAGCCTGGGAGGGCACTTACTGGCAACAGCAGTGTCTATTTGTGCAGTACAAAATATATGGACAGAGACAATGGAAAAATAAGAAGACTGTTGGGTACTCAAGAACTCTCTGTGACCTTACCAGTGAAACCTCAGACATACAGGAACCTTATTACGGGAGGGTGAGGGCGGCCTCGGCTGGGAGCTACTCAGAATGGAGCATGACGCCGCGTTCACTCCCTGGTGGGAAAGAGCAAAAGGTTTA</p> <p>ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA</p>
Protein Sequence:	<p>>RC215420 representing NM_181310</p> <p>Red=Cloning site Green=Tags(s)</p> <p>MMPKHCFGLGLISFFLTGVAGTQSTHESLKPQRVQFQSRNFHNLQWQGRALTNSSVYFVQYKIYGQRQWKNKEDCWGTQELSCDLTSETSDIQEPYGRVRAASAGSYSEWSMTPRFTPWWERAKGL</p> <p>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Restriction Sites:	Sgfl-MluI


[View online »](#)

Cloning Scheme:



ACCN: NM_181310

ORF Size: 390 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_181310.2</u>
RefSeq Size:	2644 bp
RefSeq ORF:	393 bp
Locus ID:	116379
UniProt ID:	<u>Q969J5</u>
Cytogenetics:	6q23.3
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway
MW:	14.9 kDa
Gene Summary:	<p>This gene encodes a member of the class II cytokine receptor family. The encoded soluble protein specifically binds to and inhibits interleukin 22 activity by blocking the interaction of interleukin 22 with its cell surface receptor. The encoded protein may be important in the regulation of inflammatory response, and has been implicated in the regulation of tumorigenesis in the colon. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2013]</p>

Product images:



Circular map for RC215420