

Product datasheet for RC215432L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TCF3 / E2A (TCF3) (NM 003200) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TCF3 / E2A (TCF3) (NM_003200) Human Tagged ORF Clone Lentiviral Particle

Symbol: TCF3 / E2A

Synonyms: AGM8; bHLHb21; E2A; E47; ITF1; p75; TCF-3; VDIR

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_003200 **ORF Size:** 1962 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC215432).

Sequence:

*anuangat

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003200.1

 RefSeq Size:
 4396 bp

 RefSeq ORF:
 1965 bp

 Locus ID:
 6929

 UniProt ID:
 P15923

Cytogenetics: 19p13.3

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Induced pluripotent stem cells, Transcription

Factors





MW:

67.4 kDa

Gene Summary:

This gene encodes a member of the E protein (class I) family of helix-loop-helix transcription factors. E proteins activate transcription by binding to regulatory E-box sequences on target genes as heterodimers or homodimers, and are inhibited by heterodimerization with inhibitor of DNA-binding (class IV) helix-loop-helix proteins. E proteins play a critical role in lymphopoiesis, and the encoded protein is required for B and T lymphocyte development. Deletion of this gene or diminished activity of the encoded protein may play a role in lymphoid malignancies. This gene is also involved in several chromosomal translocations that are associated with lymphoid malignancies including pre-B-cell acute lymphoblastic leukemia (t(1;19), with PBX1), childhood leukemia (t(19;19), with TFPT) and acute leukemia (t(12;19), with ZNF384). Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the short arm of chromosome 9. [provided by RefSeq, Sep 2011]