

## Product datasheet for RC215435L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## GAB2 (NM\_080491) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** GAB2 (NM\_080491) Human Tagged ORF Clone Lentiviral Particle

Symbol: GAB2 **Mammalian Cell** 

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

mGFP Tag:

ACCN: NM\_080491 **ORF Size:** 2028 bp

**ORF Nucleotide** 

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC215435).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 080491.1

RefSeq Size: 6007 bp RefSeq ORF: 2031 bp Locus ID: 9846 **UniProt ID:** Q9UQC2 **Cytogenetics:** 11q14.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Chronic myeloid leukemia, Fc epsilon RI signaling pathway, Fc gamma R-mediated

phagocytosis







MW:

74.3 kDa

**Gene Summary:** 

This gene is a member of the GRB2-associated binding protein (GAB) gene family. These proteins contain pleckstrin homology (PH) domain, and bind SHP2 tyrosine phosphatase and GRB2 adapter protein. They act as adapters for transmitting various signals in response to stimuli through cytokine and growth factor receptors, and T- and B-cell antigen receptors. The protein encoded by this gene is the principal activator of phosphatidylinositol-3 kinase in response to activation of the high affinity IgE receptor. Two alternatively spliced transcripts encoding different isoforms have been described for this gene. [provided by RefSeq, Nov 2009]