

Product datasheet for RC215442L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Syntenin (SDCBP) (NM_001007068) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Syntenin (SDCBP) (NM_001007068) Human Tagged ORF Clone Lentiviral Particle

Symbol: Syntenin

Synonyms: MDA-9; MDA9; ST1; SYCL; TACIP18

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001007068

ORF Size: 876 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC215442).

OTI Disclaimer:

Cytogenetics:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001007068.1, NP 001007069.1</u>

 RefSeq Size:
 2155 bp

 RefSeq ORF:
 879 bp

 Locus ID:
 6386

 UniProt ID:
 000560

Protein Families: Druggable Genome, Transmembrane

8q12.1

MW: 31.6 kDa





Gene Summary:

The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskeleton. The syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. Related pseudogenes have been identified on multiple chromosomes. [provided by RefSeq, Jan 2017]