

# Product datasheet for RC215533L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## FGFR3 (NM 000142) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** FGFR3 (NM\_000142) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ACH; CD333; CEK2; HSFGFR3EX; JTK4 Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 000142 ACCN: **ORF Size:** 

**ORF Nucleotide** 

2418 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC215533).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000142.2

RefSeq Size: 4093 bp RefSeq ORF: 2421 bp Locus ID: 2261 **UniProt ID:** P22607 Cytogenetics: 4p16.3

**Domains:** pkinase, TyrKc, S\_TKc, ig, IGc2, IG

**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane





#### FGFR3 (NM\_000142) Human Tagged ORF Clone Lentiviral Particle - RC215533L3V

**Protein Pathways:** Bladder cancer, Endocytosis, MAPK signaling pathway, Pathways in cancer, Regulation of actin

cytoskeleton

**MW:** 87.71 kDa

**Gene Summary:** This gene encodes a member of the fibroblast growth factor receptor (FGFR) family, with its

amino acid sequence being highly conserved between members and among divergent species. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds acidic and basic fibroblast growth hormone and plays a role in bone development and maintenance. Mutations in this gene lead to craniosynostosis and multiple types of skeletal

dysplasia. [provided by RefSeq, Aug 2017]