

Product datasheet for RC215628L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EDA (NM_001005610) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: EDA (NM_001005610) Human Tagged ORF Clone Lentiviral Particle

Symbol: EDA

Synonyms: ECTD1; ED1-A1; ED1-A2; EDA-A1; EDA-A2; EDA1; EDA2; HED; HED1; ODT1; STHAGX1;

TNLG7C; XHED; XLHED

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001005610

ORF Size: 534 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC215628).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001005610.1</u>, <u>NP 001005610.1</u>

RefSeq Size: 1089 bp
RefSeq ORF: 408 bp
Locus ID: 1896
UniProt ID: Q92838

Cytogenetics: Xq13.1

Protein Families: Druggable Genome, Secreted Protein, Transmembrane





EDA (NM_001005610) Human Tagged ORF Clone Lentiviral Particle - RC215628L3V

Protein Pathways: Cytokine-cytokine receptor interaction

MW: 19 kDa

Gene Summary: The protein encoded by this gene is a type II membrane protein that can be cleaved by furin

to produce a secreted form. The encoded protein, which belongs to the tumor necrosis factor

family, acts as a homotrimer and may be involved in cell-cell signaling during the

development of ectodermal organs. Defects in this gene are a cause of ectodermal dysplasia,

anhidrotic, which is also known as X-linked hypohidrotic ectodermal dysplasia. Several transcript variants encoding many different isoforms have been found for this gene.

[provided by RefSeq, Jul 2008]