

## Product datasheet for RC215661L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MDA5 (IFIH1) (NM\_022168) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** MDA5 (IFIH1) (NM\_022168) Human Tagged ORF Clone Lentiviral Particle

Symbol: MDA5

Synonyms: AGS7; Hlcd; IDDM19; MDA-5; MDA5; RLR-2; SGMRT1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_022168

ORF Size: 3075 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC215661).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 022168.2

 RefSeq Size:
 3434 bp

 RefSeq ORF:
 3078 bp

 Locus ID:
 64135

 UniProt ID:
 Q9BYX4

 Cytogenetics:
 2q24.2

**Domains:** DEAD, helicase\_C

**Protein Pathways:** RIG-I-like receptor signaling pathway





MW:

116.5 kDa

**Gene Summary:** 

IFIH1 encodes MDA5 which is an intracellular sensor of viral RNA that triggers the innate immune response. Sensing RNA length and secondary structure, MDA5 binds dsRNA oligonucleotides with a modified DExD/H-box helicase core and a C-terminal domain, thus leading to a proinflammatory response that includes interferons. It has been shown that Coronaviruses (CoVs) as well as various other virus families, are capable of evading the MDA5-dependent interferon response, thus impeding the activation of the innate immune response to infection. MDA5 has also been shown to play an important role in enhancing natural killer cell function in malaria infection. In addition to its protective role in antiviral responses, MDA5 has been implicated in autoimmune and autoinflammatory diseases such as type 1 diabetes, systemic lupus erythematosus, and Aicardi-Goutieres syndrome[provided by RefSeq, Jul 2020]