

Product datasheet for RC215704L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PPM1B (NM_001033557) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PPM1B (NM_001033557) Human Tagged ORF Clone Lentiviral Particle

Symbol: PPM1B

Synonyms: PP2C-beta; PP2C-beta-X; PP2CBETA; PPC2BETAX

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001033557

ORF Size: 1140 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC215704).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001033557.1</u>

 RefSeq Size:
 3056 bp

 RefSeq ORF:
 1143 bp

 Locus ID:
 5495

 UniProt ID:
 075688

Cytogenetics: 2p21

Protein Families: Druggable Genome, Phosphatase, Stem cell - Pluripotency

Protein Pathways: MAPK signaling pathway





ORIGENE

MW: 41.9 kDa

Gene Summary: The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein

phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. This phosphatase has been shown to dephosphorylate cyclin-dependent kinases (CDKs), and thus may be involved in cell cycle control. Overexpression of this phosphatase is reported to cause cell-growth arrest or cell death. Alternative splicing results

in multiple transcript variants encoding different isoforms. Additional transcript variants have been described, but currently do not represent full-length sequences. [provided by RefSeq, Jul

20081