

Product datasheet for RC215712L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HIST1H4G (H4C7) (NM_003547) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: HIST1H4G (H4C7) (NM_003547) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: H4C7

Synonyms: H4/l; H4FL; HIST1H4G

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Sgfl-Mlul

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC215712).

Sequence:

Restriction Sites: Cloning Scheme:

Cloning sites used for ORF Shuttling:

Sgf 1 ORF Mlu 1

GGG ATC GC ATG ---//--- NNN ACG CGT ---



 $[\]ensuremath{^*}$ The last codon before the Stop codon of the ORF.

ACCN: NM_003547

ORF Size: 294 bp





HIST1H4G (H4C7) (NM_003547) Human Tagged Lenti ORF Clone - RC215712L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 003547.2</u>

RefSeq Size: 367 bp
RefSeq ORF: 297 bp
Locus ID: 8369
UniProt ID: Q99525

Protein Pathways: Systemic lupus erythematosus

6p22.2

MW: 11 kDa

Cytogenetics:

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene

cluster on chromosome 6. [provided by RefSeq, Aug 2015]