

Product datasheet for RC215735L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PKN1 (NM_213560) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PKN1 (NM_213560) Human Tagged ORF Clone Lentiviral Particle

Symbol: PKN1

Synonyms: DBK; PAK-1; PAK1; PKN; PKN-ALPHA; PRK1; PRKCL1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK ACCN: NM_213560

ORF Size: 2844 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC215735).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 213560.1, NP 998725.1

 RefSeq Size:
 2964 bp

 RefSeq ORF:
 2847 bp

 Locus ID:
 5585

 UniProt ID:
 Q16512

 Cytogenetics:
 19p13.12

Protein Families: Druggable Genome, Protein Kinase

MW: 104.5 kDa







Gene Summary:

The protein encoded by this gene belongs to the protein kinase C superfamily. This kinase is activated by Rho family of small G proteins and may mediate the Rho-dependent signaling pathway. This kinase can be activated by phospholipids and by limited proteolysis. The 3-phosphoinositide dependent protein kinase-1 (PDPK1/PDK1) is reported to phosphorylate this kinase, which may mediate insulin signals to the actin cytoskeleton. The proteolytic activation of this kinase by caspase-3 or related proteases during apoptosis suggests its role in signal transduction related to apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]