

Product datasheet for RC215836L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STAT3 (NM_139276) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: STAT3 (NM_139276) Human Tagged ORF Clone Lentiviral Particle

Symbol: STAT3

Synonyms: ADMIO; ADMIO1; APRF; HIES

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_139276

 ORF Size:
 2310 bp

ORF Nucleotide

OTI Disclaimer:

otido Th

Sequence:

Domains:

The ORF insert of this clone is exactly the same as (RC215836).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 139276.2</u>

 RefSeq Size:
 4978 bp

 RefSeq ORF:
 2313 bp

 Locus ID:
 6774

 UniProt ID:
 P40763

 Cytogenetics:
 17q21.2

Protein Families: Druggable Genome, Transcription Factors

SH2, STAT





STAT3 (NM_139276) Human Tagged ORF Clone Lentiviral Particle - RC215836L3V

Protein Pathways: Acute myeloid leukemia, Adipocytokine signaling pathway, Chemokine signaling pathway, Jak-

STAT signaling pathway, Pancreatic cancer, Pathways in cancer

MW: 87.9 kDa

Gene Summary: The protein encoded by this gene is a member of the STAT protein family. In response to

cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. This protein mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of this protein. PIAS3 protein is a specific inhibitor of this protein. This gene also plays a role in regulating host response to viral and bacterial infections. Mutations in this gene are associated with infantile-onset multisystem autoimmune disease and hyper-immunoglobulin E syndrome. [provided

by RefSeq, Aug 2020]