

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC215863L4V

C1orf106 (INAVA) (NM_018265) Human Tagged ORF Clone Lentiviral Particle

Product data:

| Product Type: | Lentiviral Particles |
|------------------------------|---|
| Product Name: | C1orf106 (INAVA) (NM_018265) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | INAVA |
| Synonyms: | C1orf106 |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-mGFP-P2A-Puro (PS100093) |
| Tag: | mGFP |
| ACCN: | NM_018265 |
| ORF Size: | 1989 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC215863). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | <u>NM 018265.1</u> |
| RefSeq Size: | 4098 bp |
| RefSeq ORF: | 1992 bp |
| Locus ID: | 55765 |
| UniProt ID: | <u>Q3KP66</u> |
| Cytogenetics: | 1q32.1 |
| MW: | 72.7 kDa |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

C1orf106 (INAVA) (NM_018265) Human Tagged ORF Clone Lentiviral Particle – RC215863L4V Clore Lentiviral Particle – RC215863L4V

Gene Summary:Expressed in peripheral macrophages and intestinal myeloid-derived cells, is required for
optimal PRR (pattern recognition receptor)-induced signaling, cytokine secretion, and
bacterial clearance. Upon stimulation of a broad range of PRRs (pattern recognition receptor)
such as NOD2 or TLR2, TLR3, TLR4, TLR5, TLR7 and TLR9, associates with YWHAQ/14-3-3T,
which in turn leads to the recruitment and activation of MAP kinases and NF-kappa-B
signaling complexes that amplifies PRR-induced downstream signals and cytokine secretion
(PubMed:28436939). In the intestine, regulates adherens junction stability by regulating the
degradation of CYTH1 and CYTH2, probably acting as substrate cofactor for SCF E3 ubiquitin-
protein ligase complexes. Stabilizes adherens junctions by limiting CYTH1-dependent ARF6
activation (PubMed:29420262).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US