

Product datasheet for RC215899L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CARD9 (NM_052813) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CARD9 (NM_052813) Human Tagged ORF Clone Lentiviral Particle

Symbol: CARD9

Synonyms: CANDF2; hCARD9

Mammalian Cell

Selection:

ORF Size:

Sequence:

None

1608 bp

Vector: pLenti-C-Myc-DDK (PS100064)

Tag:Myc-DDKACCN:NM_052813

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC215899).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 052813.2

 RefSeq Size:
 2136 bp

 RefSeq ORF:
 1611 bp

 Locus ID:
 64170

 UniProt ID:
 Q9H257

 Cytogenetics:
 9q34.3

Protein Families: Druggable Genome

Protein Pathways: NOD-like receptor signaling pathway





ORIGENE

MW: 62.1 kDa

Gene Summary:

The protein encoded by this gene is a member of the CARD protein family, which is defined by the presence of a characteristic caspase-associated recruitment domain (CARD). CARD is a protein interaction domain known to participate in activation or suppression of CARD containing members of the caspase family, and thus plays an important regulatory role in cell apoptosis. This protein was identified by its selective association with the CARD domain of BCL10, a postive regulator of apoptosis and NF-kappaB activation, and is thought to function as a molecular scaffold for the assembly of a BCL10 signaling complex that activates NF-kappaB. Several alternatively spliced transcript variants have been observed, but their full-length nature is not clearly defined. [provided by RefSeq, Jul 2008]