

### Product datasheet for RC215900L4

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

OriGene Technologies, Inc.

# CN: techsupport@origene.cn

## PREX2 (NM\_025170) Human Tagged Lenti ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: PREX2 (NM\_025170) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: PREX2

Synonyms: DEP.2; DEPDC2; P-REX2; PPP1R129

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC215900).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_025170

ORF Size: 2937 bp





### PREX2 (NM\_025170) Human Tagged Lenti ORF Clone - RC215900L4

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 025170.4</u>

 RefSeq Size:
 3024 bp

 RefSeq ORF:
 2940 bp

 Locus ID:
 80243

 UniProt ID:
 Q70Z35

 Cytogenetics:
 8q13.2

**MW:** 111.9 kDa

**Gene Summary:** The protein encoded by this gene belongs to the phosphatidylinositol 3,4,5-trisphosphate


(PIP3)-dependent Rac exchanger (PREX) family, which are Dbl-type guanine-nucleotide exchange factors for Rac family small G proteins. Structural domains of this protein include the catalytic diffuse B-cell lymphoma homology and pleckstrin homology (DHPH) domain, two disheveled, EGL-10, and pleckstrin homology (DEP) domains, two PDZ domains, and a C-terminal inositol polyphosphate-4 phosphatase (IP4P) domain that is found in one of the isoforms. This protein facilitates the exchange of GDP for GTP on Rac1, allowing the GTP-bound Rac1 to activate downstream effectors. Studies also show that the pleckstrin homology domain of this protein interacts with the phosphatase and tensin homolog (PTEN) gene product to inhibit PTEN phosphatase activity, thus activating the phosphoinositide-3 kinase (PI3K) signaling pathway. Conversely, the PTEN gene product has also been shown to inhibit the GEF activity of this protein. This gene plays a role in insulin-signaling pathways, and either

mutations or overexpression of this gene have been observed in some cancers. [provided by

RefSeq, Apr 2016]



# **Product images:**



Circular map for RC215900L4