

Product datasheet for RC215948L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CREM (NM_001881) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CREM (NM_001881) Human Tagged ORF Clone Lentiviral Particle

Symbol: CREM

Synonyms: CREM-2; hCREM-2; ICER

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001881

ORF Size: 411 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC215948).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001881.3, NP 001872.3

 RefSeq Size:
 1304 bp

 RefSeq ORF:
 414 bp

 Locus ID:
 1390

 UniProt ID:
 Q03060

Cytogenetics: 10p11.21

Domains: pKID, BRLZ

Protein Families: Druggable Genome, Transcription Factors





ORIGENE

MW: 14.9 kDa

Gene Summary:

This gene encodes a bZIP transcription factor that binds to the cAMP responsive element found in many viral and cellular promoters. It is an important component of cAMP-mediated signal transduction during the spermatogenetic cycle, as well as other complex processes. Alternative promoter and translation initiation site usage allows this gene to exert spatial and temporal specificity to cAMP responsiveness. Multiple alternatively spliced transcript variants encoding several different isoforms have been found for this gene, with some of them functioning as activators and some as repressors of transcription. [provided by RefSeq, Jul 2008]