

Product datasheet for RC215957L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UGT1A6 (NM_001072) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: UGT1A6 (NM_001072) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: UGT1A6

Synonyms: GNT1; HLUGP; HLUGP1; hUG-BR1; UDPGT 1-6; UGT-1A; UGT-1C; UGT-1E; UGT-1F;

UGT1; UGT1-01; UGT1-03; UGT1-05; UGT1-06; UGT1.1; UGT1.3; UGT1.5; UGT1.6; UGT1A;

UGT1A1; UGT1A3; UGT1A5; UGT1A6S; UGT1C; UGT1E; UGT1F

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide


Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:



The ORF insert of this clone is exactly the same as(RC215957).



^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001072



UGT1A6 (NM_001072) Human Tagged Lenti ORF Clone - RC215957L1

ORF Size: 1596 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

UDPGT

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001072.2</u>

 RefSeq Size:
 2435 bp

 RefSeq ORF:
 1599 bp

 Locus ID:
 54578

 UniProt ID:
 P19224

 Cytogenetics:
 2q37.1

Domains:

Protein Families: Transmembrane

Protein Pathways: Androgen and estrogen metabolism, Ascorbate and aldarate metabolism, Drug metabolism -

cytochrome P450, Drug metabolism - other enzymes, Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Pentose and glucuronate interconversions, Porphyrin and

chlorophyll metabolism, Retinol metabolism, Starch and sucrose metabolism


MW: 60.75 kDa




Gene Summary:

This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. The enzyme encoded by this gene is active on phenolic and planar compounds. Alternative splicing in the unique 5' end of this gene results in two transcript variants. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RC215957L1



Double digestion of RC215957L1 using Sgfl and Mul