

## Product datasheet for RC215965L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SAMD8 (NM\_144660) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type:** Lentiviral Particles

Product Name: SAMD8 (NM 144660) Human Tagged ORF Clone Lentiviral Particle

Symbol: SAMD8

Synonyms: HEL-177; SMSr

**Mammalian Cell** 

Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_144660

ORF Size: 978 bp

**ORF Nucleotide** 

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC215965).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 144660.2

RefSeq Size: 7009 bp
RefSeq ORF: 981 bp
Locus ID: 142891
UniProt ID: Q96LT4
Cytogenetics: 10q22.2
Domains: SAM

**Protein Families:** Transmembrane





ORIGENE

**MW:** 37.6 kDa

**Gene Summary:** Sphingomyelin synthases synthesize sphingolipids through transfer of a phosphatidyl head

group on to the primary hydroxyl of ceramide. SAMD8 is an endoplasmic reticulum (ER)

transferase that has no sphingomyelin synthase activity but can convert

phosphatidylethanolamine (PE) and ceramide to ceramide phosphoethanolamine (CPE) albeit

with low product yield. Appears to operate as a ceramide sensor to control ceramide

homeostasis in the endoplasmic reticulum rather than a converter of ceramides. Seems to be

critical for the integrity of the early secretory pathway.[UniProtKB/Swiss-Prot Function]