

## **Product datasheet for RC215973**

# COX7A2 (NM 001865) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** COX7A2 (NM\_001865) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: COX7A2

Synonyms: COX7AL; COX7AL1; COXVIIa-L; COXVIIAL; VIIAL

Mammalian Cell Neomycin

Selection:

**ORF Nucleotide** 

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

>RC215973 ORF sequence

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCTGTGGAATCTGCTGGCTCTTCGTCAGATTGGGCAGAGGACGATAAGCACTGCTTCCCGCAGGCATT TTAAAAATAAAGTTCCGGAGAAGCAAAAACTGTTCCAGGAGGATGATGAAATTCCACTGTATCTAAAGGG TGGGGTAGCTGATGCCCTCCTGTATAGAGCCACCATGATTCTTACAGTTGGTAGAACAGCATATGCCATA

TATGAGCTGGCTGTGGCTTCATTTCCCAAGAAGCAGGAG

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAG**GTTTAA** 

**Protein Sequence:** >RC215973 protein sequence

Red=Cloning site Green=Tags(s)

MLWNLLALRQIGQRTISTASRRHFKNKVPEKQKLFQEDDEIPLYLKGGVADALLYRATMILTVGGTAYAI

YELAVASFPKKQE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6449">https://cdn.origene.com/chromatograms/mk6449</a> c04.zip

Restriction Sites: Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_001865

ORF Size: 249 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001865.4</u>

RefSeq Size: 708 bp RefSeq ORF: 252 bp Locus ID: 1347

### COX7A2 (NM\_001865) Human Tagged ORF Clone - RC215973

**UniProt ID:** P14406 Cytogenetics: 6q14.1 **Domains:** COX7a

**Protein Families:** Transmembrane

**Protein Pathways:** Alzheimer's disease, Cardiac muscle contraction, Huntington's disease, Oxidative

phosphorylation, Parkinson's disease


MW: 9.4 kDa

**Gene Summary:** Cytochrome c oxidase, the terminal component of the mitochondrial respiratory chain,


catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of three catalytic subunits encoded by mitochondrial genes, and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, while the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (liver isoform) of subunit VIIa, with this polypeptide being present in both muscle and non-muscle tissues. In addition to polypeptide 2, subunit VIIa includes polypeptide 1 (muscle isoform), which is present only in muscle tissues, and a related protein, which is present in all tissues. Alternative splicing results in multiple transcript variants. Related pseudogenes have been

identified on chromosomes 4 and 14. [provided by RefSeq, Oct 2009]

## **Product images:**



Circular map for RC215973



Western blot validation of overexpression lysate (Cat# [LY419693]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC215973 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).