

Product datasheet for RC216073L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Junctional Adhesion Molecule C (JAM3) (NM 032801) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Junctional Adhesion Molecule C (JAM3) (NM_032801) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Junctional Adhesion Molecule C

Synonyms: JAM-2; JAM-3; JAM-C; JAMC

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC216073).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_032801

ORF Size: 1065 bp





Junctional Adhesion Molecule C (JAM3) (NM_032801) Human Tagged Lenti ORF Clone – RC216073L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 032801.3</u>, <u>NP 116190.2</u>

 RefSeq Size:
 3675 bp

 RefSeq ORF:
 933 bp

 Locus ID:
 83700

 UniProt ID:
 Q9BX67

 Cytogenetics:
 11q25

Domains: ig, IGv, IGc2, IG

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Epithelial cell signaling in Helicobacter pylori infection,

Leukocyte transendothelial migration, Tight junction

MW: 36.5 kDa

Gene Summary: Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell


sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. The protein encoded by this immunoglobulin superfamily gene member is localized in the tight junctions between high endothelial cells. Unlike other proteins in this family, the this protein is unable to adhere to leukocyte cell lines and only forms weak homotypic interactions. The encoded protein is a member of the junctional adhesion molecule protein family and acts as a receptor for

hemorrhagic destruction of the brain, subependymal calcification, and congenital cataracts. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Apr 2011]

another member of this family. A mutation in an intron of this gene is associated with



Product images:



Circular map for RC216073L3