

## Product datasheet for RC216080L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## P Glycoprotein (ABCB1) (NM\_000927) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: P Glycoprotein (ABCB1) (NM 000927) Human Tagged ORF Clone Lentiviral Particle

Symbol: P Glycoprotein

Synonyms: ABC20; CD243; CLCS; GP170; MDR1; p-170; P-GP; PGY1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_000927 **ORF Size:** 3840 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216080).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000927.3

 RefSeq Size:
 4872 bp

 RefSeq ORF:
 3843 bp

 Locus ID:
 5243

 UniProt ID:
 P08183

 Cytogenetics:
 7q21.12

**Domains:** ABC\_membrane, ABC\_tran, AAA

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane





**Protein Pathways:** ABC transporters

MW: 141.3 kDa

**Gene Summary:** The membrane-associated protein encoded by this gene is a member of the superfamily of

ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance. The protein encoded by this gene is an ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier.

to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier. Mutations in this gene are associated with colchicine resistance and Inflammatory bowel disease 13. Alternative splicing and the use of alternative promoters results in multiple

transcript variants. [provided by RefSeq, Feb 2017]