

Product datasheet for RC216116L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Syntenin (SDCBP) (NM 001007069) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Syntenin (SDCBP) (NM 001007069) Human Tagged ORF Clone Lentiviral Particle

Symbol: SDCBF

Synonyms: MDA-9; MDA9; ST1; SYCL; TACIP18

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001007069

ORF Size: 891 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC216116).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001007069.1</u>, <u>NP 001007070.1</u>

 RefSeq Size:
 2170 bp

 RefSeq ORF:
 894 bp

 Locus ID:
 6386

 UniProt ID:
 000560

 Cytogenetics:
 8q12.1

Protein Families: Druggable Genome, Transmembrane

MW: 32.3 kDa





Gene Summary:

The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskeleton. The syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. Related pseudogenes have been identified on multiple chromosomes. [provided by RefSeq, Jan 2017]