

## Product datasheet for RC216130L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MEKK3 (MAP3K3) (NM\_203351) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: MEKK3 (MAP3K3) (NM 203351) Human Tagged ORF Clone Lentiviral Particle

Symbol: MEKK3

Synonyms: MAPKKK3; MEKK3

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_203351 **ORF Size:** 1971 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216130).

OTI Disclaimer:

Sequence:

mer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 203351.1

 RefSeq Size:
 4843 bp

 RefSeq ORF:
 1974 bp

 Locus ID:
 4215

 UniProt ID:
 Q99759

 Cytogenetics:
 17q23.3

**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** GnRH signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway





ORIGENE

**MW:** 73.9 kDa

**Gene Summary:** 

This gene product is a 626-amino acid polypeptide that is 96.5% identical to mouse Mekk3. Its catalytic domain is closely related to those of several other kinases, including mouse Mekk2, tobacco NPK, and yeast Ste11. Northern blot analysis revealed a 4.6-kb transcript that appears to be ubiquitously expressed. This protein directly regulates the stress-activated protein kinase (SAPK) and extracellular signal-regulated protein kinase (ERK) pathways by activating SEK and MEK1/2 respectively; it does not regulate the p38 pathway. In cotransfection assays, it enhanced transcription from a nuclear factor kappa-B (NFKB)-dependent reporter gene, consistent with a role in the SAPK pathway. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]