

Product datasheet for RC216188L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Shugoshin (SGO1) (NM_001012413) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Shugoshin (SGO1) (NM_001012413) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: Shugoshin

Synonyms: CAID; NY-BR-85; SGO; SGOL1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clo


Sequence:

The ORF insert of this clone is exactly the same as(RC216188).

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_001012413

ORF Size: 774 bp



Shugoshin (SGO1) (NM_001012413) Human Tagged Lenti ORF Clone - RC216188L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001012413.1</u>, <u>NP 001012413.1</u>

 RefSeq Size:
 1149 bp

 RefSeq ORF:
 777 bp

 Locus ID:
 151648

 UniProt ID:
 Q5FBB7

 Cytogenetics:
 3p24.3

Protein Pathways: Oocyte meiosis

MW: 29.3 kDa

Gene Summary: The protein encoded by this gene is a member of the shugoshin family of proteins. This


protein is thought to protect centromeric cohesin from cleavage during mitotic prophase by preventing phosphorylation of a cohesin subunit. Reduced expression of this gene leads to the premature loss of centromeric cohesion, mis-segregation of sister chromatids, and mitotic arrest. Evidence suggests that this protein also protects a small subset of cohesin found along the length of the chromosome arms during mitotic prophase. An isoform lacking exon 6 has been shown to play a role in the cohesion of centrioles (PMID: 16582621 and PMID:18331714). Mutations in this gene have been associated with Chronic Atrial and Intestinal Dysrhythmia (CAID) syndrome, characterized by the co-occurrence of Sick Sinus Syndrome (SSS) and Chronic Intestinal Pseudo-obstruction (CIPO) within the first four decades of life (PMID:25282101). Fibroblast cells from CAID patients exhibited both increased cell proliferation and higher rates of senescence. Pseudogenes of this gene have been found on

chromosomes 1 and 7. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Mar 2015]



Product images:



Circular map for RC216188L3