

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC216209L3V

TLR10 (NM_001017388) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	TLR10 (NM_001017388) Human Tagged ORF Clone Lentiviral Particle
Symbol:	TLR10
Synonyms:	CD290
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001017388
ORF Size:	2433 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC216209).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001017388.1, NP 001017388.1</u>
RefSeq Size:	3484 bp
RefSeq ORF:	2436 bp
Locus ID:	81793
UniProt ID:	<u>Q9BXR5</u>
Cytogenetics:	4p14
Protein Families:	Druggable Genome, Transmembrane
MW:	94.6 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which
plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are
highly conserved from Drosophila to humans and share structural and functional similarities.
They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on
infectious agents, and mediate the production of cytokines necessary for the development of
effective immunity. The various TLRs exhibit different patterns of expression. This gene is
most highly expressed in lymphoid tissues such as spleen, lymph node, thymus, and tonsil.
Multiple alternatively spliced transcript variants which encode different protein isoforms have
been found for this gene. [provided by RefSeq, Aug 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US