

## Product datasheet for RC216427L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CYP4F2 (NM\_001082) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** CYP4F2 (NM\_001082) Human Tagged ORF Clone Lentiviral Particle

Symbol: CYP4F2
Synonyms: CPF2

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001082 **ORF Size:** 1560 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216427).

Sequence:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001082.3

 RefSeq Size:
 2360 bp

 RefSeq ORF:
 1563 bp

 Locus ID:
 8529

 UniProt ID:
 P78329

 Cytogenetics:
 19p13.12

**Domains:** p450

**Protein Families:** Druggable Genome, P450, Transmembrane





## CYP4F2 (NM\_001082) Human Tagged ORF Clone Lentiviral Particle - RC216427L4V

**Protein Pathways:** Arachidonic acid metabolism, Metabolic pathways

**MW:** 59.7 kDa

**Gene Summary:** This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum. The enzyme starts the process of inactivating and degrading

leukotriene B4, a potent mediator of inflammation. This gene is part of a cluster of cytochrome P450 genes on chromosome 19. Another member of this family, CYP4F11, is

approximately 16 kb away. [provided by RefSeq, Jul 2008]