

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC216461L4V

MTMR3 (NM_153050) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	MTMR3 (NM_153050) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MTMR3
Synonyms:	FYVE-DSP1; ZFYVE10
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_153050
ORF Size:	3510 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC216461).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 153050.2, NP 694690.1</u>
RefSeq Size:	8924 bp
RefSeq ORF:	3513 bp
Locus ID:	8897
UniProt ID:	<u>Q13615</u>
Cytogenetics:	22q12.2
Protein Families:	Druggable Genome, Phosphatase
MW:	130.6 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a member of the myotubularin dual specificity protein phosphatase gene
family. The encoded protein is structurally similar to myotubularin but in addition contains a
FYVE domain and an N-terminal PH-GRAM domain. The protein can self-associate and also
form heteromers with another myotubularin related protein. The protein binds to
phosphoinositide lipids through the PH-GRAM domain, and can hydrolyze
phosphatidylinositol(3)-phosphate and phosphatidylinositol(3,5)-biphosphate in vitro. The
encoded protein has been observed to have a perinuclear, possibly membrane-bound,
distribution in cells, but it has also been found free in the cytoplasm. Multiple transcript
variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul
2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US