

## Product datasheet for RC216463L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MAP3K12 (NM\_006301) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** MAP3K12 (NM\_006301) Human Tagged ORF Clone Lentiviral Particle

Symbol: MAP3K12

Synonyms: DLK; MEKK12; MUK; ZPK; ZPKP1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_006301

 ORF Size:
 2577 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC216463).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 006301.3

 RefSeq Size:
 3541 bp

 RefSeq ORF:
 2580 bp

 Locus ID:
 7786

 UniProt ID:
 Q12852

 Cytogenetics:
 12q13.13

**Domains:** pkinase, TyrKc, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase





## MAP3K12 (NM\_006301) Human Tagged ORF Clone Lentiviral Particle - RC216463L3V

**Protein Pathways:** MAPK signaling pathway

**MW:** 93.7 kDa

**Gene Summary:** This gene encodes a member of the serine/threonine protein kinase family. This kinase

contains a leucine-zipper domain and is predominately expressed in neuronal cells. The phosphorylation state of this kinase in synaptic terminals was shown to be regulated by membrane depolarization via calcineurin. This kinase forms heterodimers with leucine zipper containing transcription factors, such as cAMP responsive element binding protein (CREB) and MYC, and thus may play a regulatory role in PKA or retinoic acid induced neuronal differentiation. Alternatively spliced transcript variants encoding different proteins have been

described.[provided by RefSeq, Jul 2010]