

## Product datasheet for RC216495L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## c Maf (MAF) (NM 005360) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** c Maf (MAF) (NM\_005360) Human Tagged ORF Clone Lentiviral Particle

Symbol:

AYGRP; c-MAF; CCA4; CTRCT21 Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 005360

**ORF Size:** 1209 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216495).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 005360.3

RefSeq Size: 2161 bp RefSeq ORF: 1212 bp Locus ID: 4094 **UniProt ID:** 075444 Cytogenetics: 16q23.2

**Domains:** bZIP Maf, BRLZ

**Protein Families:** Druggable Genome, Transcription Factors





ORIGENE

**MW:** 41.8 kDa

**Gene Summary:** 

The protein encoded by this gene is a DNA-binding, leucine zipper-containing transcription factor that acts as a homodimer or as a heterodimer. Depending on the binding site and binding partner, the encoded protein can be a transcriptional activator or repressor. This protein plays a role in the regulation of several cellular processes, including embryonic lens fiber cell development, increased T-cell susceptibility to apoptosis, and chondrocyte terminal differentiation. Defects in this gene are a cause of juvenile-onset pulverulent cataract as well as congenital cerulean cataract 4 (CCA4). Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2010]