

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC216523L4V

CCDC50 (NM_178335) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	CCDC50 (NM_178335) Human Tagged ORF Clone Lentiviral Particle
Symbol:	CCDC50
Synonyms:	C3orf6; DFNA44; YMER
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_178335
ORF Size:	1446 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC216523).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 178335.1</u>
RefSeq Size:	2454 bp
RefSeq ORF:	1449 bp
Locus ID:	152137
UniProt ID:	<u>Q8IVM0</u>
Cytogenetics:	3q28
MW:	56.2 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a soluble, cytoplasmic, tyrosine-phosphorylated protein with multiple
ubiquitin-interacting domains. Mutations in this gene cause nonsyndromic, postlingual,
progressive sensorineural DFNA44 hearing loss. In mouse, the protein is expressed in the
inner ear during development and postnatal maturation and associates with microtubule-
based structures. This protein may also function as a negative regulator of NF-kB signaling
and as an effector of epidermal growth factor (EGF)-mediated cell signaling. Alternative
splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq,
Oct 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US