

Product datasheet for RC216634L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hexokinase 1 (HK1) (NM 033497) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Hexokinase 1 (HK1) (NM_033497) Human Tagged ORF Clone Lentiviral Particle

Symbol: HK'

Synonyms: hexokinase; HK; HK1-ta; HK1-tb; HK1-tc; HKD; HKI; HMSNR; HXK1; NEDVIBA; RP79

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM_033497

ORF Size: 2763 bp

ORF Nucleotide Sequence:

The ORF insert of this clone is exactly the same as(RC216634).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 033497.2, NP 277032.1

 RefSeq Size:
 3832 bp

 RefSeq ORF:
 2766 bp

 Locus ID:
 3098

 UniProt ID:
 P19367

 Cytogenetics:
 10q22.1

Protein Families: Druggable Genome





Hexokinase 1 (HK1) (NM_033497) Human Tagged ORF Clone Lentiviral Particle - RC216634L3V

Protein Pathways: Amino sugar and nucleotide sugar metabolism, Fructose and mannose metabolism,

Galactose metabolism, Glycolysis / Gluconeogenesis, Insulin signaling pathway, Metabolic

pathways, Starch and sucrose metabolism, Type II diabetes mellitus

MW: 102.7 kDa

Gene Summary: Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most

glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in several transcript variants which encode different isoforms, some of which

are tissue-specific. [provided by RefSeq, Apr 2016]