

## Product datasheet for **RC216645**

### Factor I (CFI) (NM\_000204) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Factor I (CFI) (NM_000204) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Factor I
Synonyms:	AHUS3; ARMD13; C3b-INA; C3BINA; FI; IF; KAF
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RC216645 representing NM\_000204  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGCATCGCC**

ATGAAGCTTCTTCATGTTTTCTGTTATTTCTGTGCTTCCACTTAAGTTTTGCAAGGTCACTTATACAT  
 CTC AAGAGGATCTGGTGGAGAAAAAGTGCCTTAGCAAAAAATATACTCACCTCTCCTGCGATAAAGTCTT  
 CTGCCAGCCATGGCAGAGATGCATTGAGGGCACCTGTGTTTGTAAACTACCGTATCAGTGCCCAAAGAAT  
 GGCAGTGCAGTGTGTGCAACTAACAGGAGAAGCTTCCCAACTACTGTCAACAAAAGAGTTTGGAAATGTC  
 TTCATCCAGGGACAAAGTTTTAAATAACGGAACATGCACAGCCGAAGGAAAGTTTAGTGTTCCTTGAA  
 GCATGGAATACAGATTCAGAGGAATAGTTGAAGTAAACTTGTGGACCAAGATAAGACAATGTTTATA  
 TGCAAAAGCAGCTGGAGCATGAGGGAAGCCAACGTGGCCTGCCTTGACCTGGGTTTCAACAAGGTGCTG  
 ATACTCAAAGAAGTTTTAAGTTGTCTGATCTCTATAAATCCACTGAATGTCTACATGTGCATTGCCG  
 AGGATTAGAGACCAGTTTGGCTGAATGTACTTTACTAAGAGAAGAACTATGGGTTACCAGGATTCGCT  
 GATGTGGTTTGTATACACAGAAAGCAGATTCTCCAATGGATGACTTCTTTCAGTGTGTGAATGGGAAAT  
 ACATTTCTCAGATGAAAGCCTGTGATGGTATCAATGATTGTGGAGACCAAGTATGAAGTGTGTTGTA  
 AGCATGCCAAGGCAAAGGCTTCCATTGCAAATCAGGTGTTTGCATTCCAAGCCAGTATCAATGCAATGGT  
 GAGGTGGACTGCATTACAGGGGAAGATGAAGTTGGCTGTGCAGGCTTTCATCTGTGGCTCAAGAAGAAA  
 CAGAAATTTGACTGCTGACATGGATGCAGAAAGAAGACGGATAAAATCATTATTACCTAAACTATCTTG  
 TGGAGTAAAAACAGAAATGCACATTGCAAGGAAACGAATTGTGGGAGGAAAGCGAGCACAACCTGGGAGAC  
 CTCCCATGGCAGGTGGCAATTAAGGATGCCAGTGAATCACCTGTGGGGGAAATTTATATTGGTGGCTGTT  
 GGATTCGACTGCTGCACATTGTCTCAGAGCCAGTAAACTCATCGTTACCAAATATGGACAACAGTAGT  
 AGACTGGATACACCCCGACCTTAAACGTATAGTAATTGAATACGTGGATAGAATATTTTCCATGAAAAAC  
 TACAATGCAGGCCTTACCAAATGACATCGCTTTGATTGAAATGAAAAAGACGAAAAACAAAAAGATT  
 GTGAGCTGCCTCGTTCCATCCCTGCCTGTGTCCCTGGTCTCCTTACCTATTCCAACCTAATGATACATG  
 CATCGTTTCTGGCTGGGACGAGAAAAAGATAACGAAAGAGTCTTTTCACTTCAGTGGGGTGAAGTTAAA  
 CTAATAAGCAACTGCTCTAAGTTTTACGGAAATCGTTTCTATGAAAAAGAAATGGAATGTGCAGGTACAT  
 ATGATGGTTCCATCGATGCCTGTAAGGGGACTCTGGAGGCCCTTAGTCTGTATGGATGCCAACAATGT  
 GACTTATGTCTGGGTGTTGTGAGTTGGGGGAAAAGTGTGAAAACAGAGTCCCAGGTGTTTACACC  
 AAAGTGGCAATATTTTACTGGATTAGCTACCATGTAGGAAGGCCTTTTATTTCTCAGTACAATGTA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:**

>RC216645 representing NM\_000204  
 Red=Cloning site Green=Tags(s)

MKLLHVFLLFLCFHLRFCKVITYTSQEDLVEKKCLAKKYTHLSCDKVFCQPWQRCIEGTCVCKLPYQCPKN  
 GTAVCATNRRSFPTYCQQKSLECLHPGKFLNNGTCTAEGKFSVSLKHGNTDSEGIVEVKLVQDKTMFI  
 CKSSWSMREANVACLDLGFQQGADTQRRFKLSDLSINSTECLHVHCRGLETSLAECTFTKRRTMGYQDFA  
 DVVYCYTQKADSPMDDFFQCVNGKYISQMKACDGINDCGDQSDDELCKKACQKGFHCKSGVCIPSQYQCNG  
 EVDCITGEDEVGCAGFASVAQEETEIL TADMDAERRRIKSLLPKLSCGVKNRMHIRRKRIVGGKRAQLGD  
 LPWQVAIKDASGITCGGIYIGGCWILTAHCLRASKTHRYQIWTTVVDWIHPDLKRIVIEYVDRIIFHEN  
 YNAGTYQNDIALIEMKKDGNKKDCELRSPACVPWSPYLFQPNDCIVSWGREGKDNERVVSLQWGEVK  
 LISNCSKFYGNRFYEKEMECAGTYDGSIDACKGDSGGPLVCM DANNVTYVWGVVSWGENCCKPEFPGVYT  
 KVANYFDWISYHVGRPFISQYNV

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:**

[https://cdn.origene.com/chromatograms/mk6482\\_e05.zip](https://cdn.origene.com/chromatograms/mk6482_e05.zip)

**Restriction Sites:**

Sgfl-Mlul

**Cloning Scheme:**


**ACCN:** NM\_000204

**ORF Size:** 1749 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
  2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
  3. Close the tube and incubate for 10 minutes at room temperature.
  4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
  5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [NM\\_000204.5](#)


**RefSeq Size:** 1963 bp

**RefSeq ORF:** 1752 bp


**Locus ID:** 3426

<b>UniProt ID:</b>	<u>P05156</u>
<b>Cytogenetics:</b>	4q25
<b>Domains:</b>	SR, Tryp_SPC, ldl_recept_a, FIMAC
<b>Protein Families:</b>	Druggable Genome, Protease, Secreted Protein
<b>Protein Pathways:</b>	Complement and coagulation cascades
<b>MW:</b>	65.72 kDa
<b>Gene Summary:</b>	This gene encodes a serine proteinase that is essential for regulating the complement cascade. The encoded preproprotein is cleaved to produce both heavy and light chains, which are linked by disulfide bonds to form a heterodimeric glycoprotein. This heterodimer can cleave and inactivate the complement components C4b and C3b, and it prevents the assembly of the C3 and C5 convertase enzymes. Defects in this gene cause complement factor I deficiency, an autosomal recessive disease associated with a susceptibility to pyogenic infections. Mutations in this gene have been associated with a predisposition to atypical hemolytic uremic syndrome, a disease characterized by acute renal failure, microangiopathic hemolytic anemia and thrombocytopenia. Primary glomerulonephritis with immune deposits and age-related macular degeneration are other conditions associated with mutations of this gene. [provided by RefSeq, Dec 2015]


### Product images:




Circular map for RC216645



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CFI (Cat# RC216645, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CFI (Cat# [TA806003]). Positive lysates [LY424865] (100ug) and [LC424865] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY424865]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC216645 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified CFI protein (Cat# [TP316645]). The protein was produced from HEK293T cells transfected with CFI cDNA clone (Cat# RC216645) using MegaTran 2.0 (Cat# [TT210002]).