

## Product datasheet for RC216665L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MASP2 (NM\_006610) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** MASP2 (NM\_006610) Human Tagged ORF Clone Lentiviral Particle

Symbol: MASP2

Synonyms: MAP-2; MAP19; MASP-2; MASP1P1; sMAP

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_006610

ORF Size: 2058 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC216665).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 006610.2</u>

 RefSeq Size:
 2460 bp

 RefSeq ORF:
 2061 bp

 Locus ID:
 10747

 UniProt ID:
 000187

 Cytogenetics:
 1p36.22

**Domains:** CCP, CUB, Tryp\_SPc, EGF\_CA, EGF

**Protein Families:** Druggable Genome, Protease, Secreted Protein





## MASP2 (NM\_006610) Human Tagged ORF Clone Lentiviral Particle - RC216665L1V

**Protein Pathways:** Complement and coagulation cascades

**MW:** 75.69 kDa

**Gene Summary:** This gene encodes a member of the peptidase S1 family of serine proteases. The encoded

preproprotein is proteolytically processed to generate A and B chains that heterodimerize to form the mature protease. This protease cleaves complement components C2 and C4 in order to generate C3 convertase in the lectin pathway of the complement system. The encoded protease also plays a role in the coagulation cascade through cleavage of prothrombin to form thrombin. Myocardial infarction and acute stroke patients exhibit reduced serum concentrations of the encoded protein. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.

[provided by RefSeq, Feb 2016]