

Product datasheet for RC216715L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PRAK (MAPKAPK5) (NM_003668) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: PRAK (MAPKAPK5) (NM_003668) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: PRAK

Synonyms: MAPKAP-K5; MK-5; MK5; PRAK

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC216715).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_003668

ORF Size: 1413 bp





PRAK (MAPKAPK5) (NM_003668) Human Tagged Lenti ORF Clone - RC216715L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 003668.2</u>

RefSeq Size:2060 bpRefSeq ORF:1416 bpLocus ID:8550

UniProt ID:

Cytogenetics: 12q24.12-q24.13

Domains: pkinase, TyrKc, S TKc

Protein Families: Druggable Genome, Protein Kinase

Q8IW41

Protein Pathways: MAPK signaling pathway


MW: 53.9 kDa

Gene Summary: The protein encoded by this gene is a tumor suppressor and member of the serine/threonine

kinase family. In response to cellular stress and proinflammatory cytokines, this kinase is activated through its phosphorylation by MAP kinases including MAPK1/ERK, MAPK14/p38-alpha, and MAPK11/p38-beta. The encoded protein is found in the nucleus but translocates to the cytoplasm upon phosphorylation and activation. This kinase phosphorylates heat shock protein HSP27 at its physiologically relevant sites. Two alternately spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Nov 2012]



Product images:



Circular map for RC216715L4