

## Product datasheet for RC216785L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL10 (NM\_000572) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** IL10 (NM\_000572) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL10

Synonyms: CSIF; GVHDS; IL-10; IL10A; TGIF

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_000572

ORF Size: 534 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216785).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000572.2

 RefSeq Size:
 1629 bp

 RefSeq ORF:
 537 bp

 Locus ID:
 3586

 UniProt ID:
 P22301

 Cytogenetics:
 1q32.1

Domains: IL10

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein



## IL10 (NM\_000572) Human Tagged ORF Clone Lentiviral Particle - RC216785L2V

**Protein Pathways:** Allograft rejection, Asthma, Autoimmune thyroid disease, Cytokine-cytokine receptor

interaction, Jak-STAT signaling pathway, Systemic lupus erythematosus, T cell receptor

signaling pathway

MW: 20.52 kDa

**Gene Summary:** The protein encoded by this gene is a cytokine produced primarily by monocytes and to a

lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid

arthritis. [provided by RefSeq, May 2020]