

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC216863L3V

ALDH8A1 (NM_170771) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	ALDH8A1 (NM_170771) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ALDH8A1
Synonyms:	ALDH12; DJ352A20.2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_170771
ORF Size:	1299 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC216863).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 170771.1</u>
RefSeq Size:	2387 bp
RefSeq ORF:	1302 bp
Locus ID:	64577
UniProt ID:	<u>Q9H2A2</u>
Cytogenetics:	6q23.3
Protein Families:	Druggable Genome
MW:	47.1 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a member of the aldehyde dehydrogenase family of proteins. The
encoded protein has been implicated in the synthesis of 9-cis-retinoic acid and in the
breakdown of the amino acid tryptophan. This enzyme converts 9-cis-retinal into the retinoid
X receptor ligand 9-cis-retinoic acid, and has approximately 40-fold higher activity with 9-cis-
retinal than with all-trans-retinal. In addition, this enzyme has been shown to catalyze the
conversion of 2-aminomuconic semialdehyde to 2-aminomuconate in the kynurenine
pathway of tryptophan catabolism. [provided by RefSeq, Jul 2018]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US