

Product datasheet for RC216887L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CLDN19 (NM 148960) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CLDN19 (NM_148960) Human Tagged ORF Clone Lentiviral Particle

Symbol: CLDN19 HOMG5 Synonyms: None

Mammalian Cell

Selection:

Vector:

pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 148960 ACCN:

ORF Size: 672 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC216887).

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer: reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 148960.1, NP 683763.1

RefSeq Size: 2859 bp RefSeq ORF: 675 bp Locus ID: 149461 **UniProt ID:** Q8N6F1 Cytogenetics: 1p34.2

Protein Families: Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Leukocyte transendothelial migration, Tight junction





ORIGENE

MW: 23 kDa

Gene Summary:

The product of this gene belongs to the claudin family. It plays a major role in tight junction-specific obliteration of the intercellular space, through calcium-independent cell-adhesion activity. Defects in this gene are the cause of hypomagnesemia renal with ocular involvement (HOMGO). HOMGO is a progressive renal disease characterized by primary renal magnesium wasting with hypomagnesemia, hypercalciuria and nephrocalcinosis associated with severe ocular abnormalities such as bilateral chorioretinal scars, macular colobomata, significant myopia and nystagmus. Alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jun 2010]