

## Product datasheet for RC216898L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IGFBP3 (NM\_001013398) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** IGFBP3 (NM\_001013398) Human Tagged ORF Clone Lentiviral Particle

Symbol: IGFBP3

Synonyms: BP-53; IBP3

Mammalian Cell Pu

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001013398

ORF Size: 891 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216898).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001013398.1</u>, <u>NP 001013416.1</u>

 RefSeq Size:
 2638 bp

 RefSeq ORF:
 894 bp

 Locus ID:
 3486

 UniProt ID:
 P17936

 Cytogenetics:
 7p12.3

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** p53 signaling pathway





ORIGENE

MW: 32.21 kDa

Gene Summary: This gene is a member of the insulin-like growth factor binding protein (IGFBP) family and

encodes a protein with an IGFBP domain and a thyroglobulin type-I domain. The protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the half-life of IGFs and altering their interaction with cell surface receptors. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

[provided by RefSeq, Jul 2008]