

## Product datasheet for RC216926L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **USP8 (NM 005154) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** USP8 (NM 005154) Human Tagged ORF Clone Lentiviral Particle

Symbol:

HumORF8; PITA4; SPG59; UBPY Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 005154 ACCN: **ORF Size:** 3354 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC216926).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 005154.2

RefSeq Size: 4250 bp RefSeq ORF: 3357 bp Locus ID: 9101 **UniProt ID:** P40818 Cytogenetics: 15q21.2

**Domains:** UCH, RHOD

**Protein Families:** Druggable Genome, Protease



## USP8 (NM\_005154) Human Tagged ORF Clone Lentiviral Particle - RC216926L1V

**Protein Pathways:** Endocytosis

MW: 127.3 kDa

**Gene Summary:** This gene encodes a protein that belongs to the ubiquitin-specific processing protease family

of proteins. The encoded protein is thought to regulate the morphology of the endosome by ubiquitination of proteins on this organelle and is involved in cargo sorting and membrane trafficking at the early endosome stage. This protein is required for the cell to enter the S phase of the cell cycle and also functions as a positive regulator in the Hedgehog signaling pathway in development. Pseudogenes of this gene are present on chromosomes 2 and 6. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]