

# Product datasheet for RC216947L1V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# IL13 (NM\_002188) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

Product Name: IL13 (NM 002188) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL13

Synonyms: IL-13; P600

Mammalian Cell None

Selection:

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 002188

ORF Size: 438 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC216947).

Sequence:

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002188.2

 RefSeq Size:
 1282 bp

 RefSeq ORF:
 441 bp

 Locus ID:
 3596

 UniProt ID:
 P35225

**Cytogenetics:** 5q31.1

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane





## IL13 (NM\_002188) Human Tagged ORF Clone Lentiviral Particle - RC216947L1V

**Protein Pathways:** Asthma, Cytokine-cytokine receptor interaction, Fc epsilon RI signaling pathway, Jak-STAT

signaling pathway

**MW:** 15.79 kDa

**Gene Summary:** This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells.

This cytokine is involved in several stages of B-cell maturation and differentiation. It upregulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-

inflammatory cytokines and chemokines. This cytokine is found to be critical to the

pathogenesis of allergen-induced asthma but operates through mechanisms independent of

IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008]