

# Product datasheet for RC217022L1V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# HCK (NM\_002110) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** HCK (NM\_002110) Human Tagged ORF Clone Lentiviral Particle

Symbol: HCK

Synonyms: JTK9; p59Hck; p61Hck

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_002110

 ORF Size:
 1578 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC217022).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002110.2

 RefSeq Size:
 2168 bp

 RefSeq ORF:
 1581 bp

 Locus ID:
 3055

 UniProt ID:
 P08631

 Cytogenetics:
 20q11.21

**Domains:** pkinase, SH2, TyrKc, SH3, S\_TKc

**Protein Families:** Druggable Genome, Protein Kinase





## HCK (NM\_002110) Human Tagged ORF Clone Lentiviral Particle - RC217022L1V

**Protein Pathways:** Chemokine signaling pathway, Fc gamma R-mediated phagocytosis

MW: 59.6 kDa

**Gene Summary:** The protein encoded by this gene is a member of the Src family of tyrosine kinases. This

protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages. It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms with different subcellular distributions are produced due to both alternative splicing and the use of alternative translation initiation codons, including a non-AUG (CUG) codon. [provided

by RefSeq, Feb 2010]