

# **Product datasheet for RC217172**

### M-CSF (CSF1) (NM 172211) Human Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** M-CSF (CSF1) (NM\_172211) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: M-CSF

Synonyms: CSF-1; MCSF

Mammalian Cell Neomycin

Selection:

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >RC217172 representing NM\_172211

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$ 

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Protein Sequence:** >RC217172 representing NM\_172211

Red=Cloning site Green=Tags(s)

MTAPGAAGRCPPTTWLGSLLLLVCLLASRSITEEVSEYCSHMIGSGHLQSLQRLIDSQMETSCQITFEFV DQEQLKDPVCYLKKAFLLVQDIMEDTMRFRDNTPNAIAIVQLQELSLRLKSCFTKDYEEHDKACVRTFYE TPLQLLEKVKNVFNETKNLLDKDWNIFSKNCNNSFAECSSQGHERQSEGSSSPQLQESVFHLLVPSVILV LLAVGGLLFYRWRRRSHQEPQRADSPLEQPEGSPLTQDDRQVELPV


**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6503">https://cdn.origene.com/chromatograms/mk6503</a> h02.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_172211

ORF Size: 768 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

#### M-CSF (CSF1) (NM\_172211) Human Tagged ORF Clone - RC217172

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 172211.3</u>

 RefSeq Size:
 1855 bp

 RefSeq ORF:
 771 bp

 Locus ID:
 1435

 UniProt ID:
 P09603

 Cytogenetics:
 1p13.3

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Hematopoietic cell lineage

**MW:** 29 kDa


**Gene Summary:** The protein encoded by this gene is a cytokine that controls the production, differentiation,

and function of macrophages. The active form of the protein is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. The encoded protein may be involved in development of the placenta. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep


2011]



## **Product images:**



Circular map for RC217172



Western blot validation of overexpression lysate (Cat# [LY406741]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC217172 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).





Coomassie blue staining of purified CSF1 protein (Cat# [TP317172]). The protein was produced from HEK293T cells transfected with CSF1 cDNA clone (Cat# RC217172) using MegaTran 2.0 (Cat# [TT210002]).