

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC217258L1V

IL32 (NM_001012718) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	IL32 (NM_001012718) Human Tagged ORF Clone Lentiviral Particle
Symbol:	IL32
Synonyms:	IL-32alpha; IL-32beta; IL-32delta; IL-32gamma; NK4; TAIF; TAIFa; TAIFb; TAIFc; TAIFd
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_001012718
ORF Size:	564 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC217258).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001012718.1</u>
RefSeq Size:	1093 bp
RefSeq ORF:	567 bp
Locus ID:	9235
UniProt ID:	<u>P24001</u>
Cytogenetics:	16p13.3
Protein Families:	Secreted Protein
MW:	21.7 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	IL32 (NM_001012718) Human Tagged ORF Clone Lentiviral Particle – RC217258L1V
Gene Summary:	This gene encodes a member of the cytokine family. The protein contains a tyrosine sulfation site, 3 potential N-myristoylation sites, multiple putative phosphorylation sites, and an RGD
	cell-attachment sequence. Expression of this protein is increased after the activation of T-cells
	by mitogens or the activation of NK cells by IL-2. This protein induces the production of

isoforms, have been characterized. [provided by RefSeq, Jul 2008]

TNFalpha from macrophage cells. Alternate transcriptional splice variants, encoding different

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US