

Product datasheet for RC217265L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Integrin alpha E (ITGAE) (NM_002208) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Integrin alpha E (ITGAE) (NM_002208) Human Tagged ORF Clone Lentiviral Particle

Symbol: Integrin alpha E
Synonyms: CD103; HUMINAE

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_002208 **ORF Size:** 3537 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC217265).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002208.4

 RefSeq Size:
 3878 bp

 RefSeq ORF:
 3540 bp

 Locus ID:
 3682

 UniProt ID:
 P38570

 Cytogenetics:
 17p13.2

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Regulation of actin cytoskeleton





Integrin alpha E (ITGAE) (NM_002208) Human Tagged ORF Clone Lentiviral Particle – RC217265L2V

MW: 130.12 kDa

Gene Summary:

Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This gene encodes an I-domain-containing alpha integrin that undergoes post-translational cleavage in the extracellular domain, yielding disulfide-linked heavy and light chains. In combination with the beta 7 integrin, this protein forms the E-cadherin binding integrin known as the human mucosal lymphocyte-1 antigen. This protein is preferentially expressed in human intestinal intraepithelial lymphocytes (IEL), and in addition to a role in adhesion, it may serve as an accessory molecule for IEL activation. [provided by RefSeq, Jul 2008]