

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC217372L4V

## TCP1 alpha (TCP1) (NM\_001008897) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	TCP1 alpha (TCP1) (NM_001008897) Human Tagged ORF Clone Lentiviral Particle
Symbol:	TCP1 alpha
Synonyms:	CCT-alpha; CCT1; CCTa; D6S230E; TCP-1-alpha
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001008897
ORF Size:	1671 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC217372).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001008897.1, NP 001008897.1</u>
RefSeq Size:	2377 bp
RefSeq ORF:	1206 bp
Locus ID:	6950
Cytogenetics:	6q25.3
MW:	60.3 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	TCP1 alpha (TCP1) (NM_001008897) Human Tagged ORF Clone Lentiviral Particle – RC217372L4V
Gene Summary:	The protein encoded by this gene is a molecular chaperone that is a member of the
	chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This
	complex consists of two identical stacked rings, each containing eight different proteins.
	Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-
	dependent manner. The complex folds various proteins, including actin and tubulin.
	Alternate transcriptional splice variants of this gene, encoding different isoforms, have been

been found. [provided by RefSeq, Jun 2010]

characterized. In addition, three pseudogenes that appear to be derived from this gene have

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US