

## **Product datasheet for RC217384L3**

# EGFR (NM\_005228) Human Tagged Lenti ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** EGFR (NM\_005228) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: EGFR

Synonyms: ERBB; ERBB1; ERRP; HER1; mENA; NISBD2; PIG61

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC217384).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





st The last codon before the Stop codon of the ORF.

**ACCN:** NM\_005228

ORF Size: 3630 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

#### EGFR (NM\_005228) Human Tagged Lenti ORF Clone - RC217384L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeq:** <u>NM 005228.3</u>, <u>NP 005219.2</u>

7p11.2

 RefSeq Size:
 5616 bp

 RefSeq ORF:
 3633 bp

 Locus ID:
 1956

 UniProt ID:
 P00533

Cytogenetics:

**Domains:** Recep L domain, pkinase, TyrKc, S TKc, Furin-like, FU

Protein Families: Adult stem cells, Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Protein

Kinase, Secreted Protein, Stem cell relevant signaling - JAK/STAT signaling pathway,

Transmembrane

**Protein Pathways:** Adherens junction, Bladder cancer, Calcium signaling pathway, Colorectal cancer, Cytokine-

cytokine receptor interaction, Dorso-ventral axis formation, Endocytosis, Endometrial cancer,

Epithelial cell signaling in Helicobacter pylori infection, ErbB signaling pathway, Focal adhesion, Gap junction, Glioma, GnRH signaling pathway, MAPK signaling pathway, Melanoma, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Prostate

cancer, Regulation of actin cytoskeleton


MW: 134.7 kDa




#### **Gene Summary:**

The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor, thus inducing receptor dimerization and tyrosine autophosphorylation leading to cell proliferation. Mutations in this gene are associated with lung cancer. EGFR is a component of the cytokine storm which contributes to a severe form of Coronavirus Disease 2019 (COVID-19) resulting from infection with severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). [provided by RefSeq, Jul 2020]

## **Product images:**



Circular map for RC217384L3



Double digestion of RC217384L3 using Sgfl and Mlul  $\,$