

## Product datasheet for **RC217402L2V**

### Metabotropic Glutamate Receptor 7 (GRM7) (NM\_000844) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Metabotropic Glutamate Receptor 7 (GRM7) (NM_000844) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Metabotropic Glutamate Receptor 7
Synonyms:	GLUR7; GPRC1G; MGLU7; MGLUR7; NEDSHBA; PPP1R87
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_000844
ORF Size:	2745 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC217402).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_000844.2</a>
RefSeq Size:	4147 bp
RefSeq ORF:	2748 bp
Locus ID:	2917
UniProt ID:	<a href="#">Q14831</a>
Cytogenetics:	3p26.1
Domains:	7tm_3, ANF_receptor



[View online »](#)

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Neuroactive ligand-receptor interaction

**MW:** 102.7 kDa

**Gene Summary:** L-glutamate is the major excitatory neurotransmitter in the central nervous system, and it activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors that have been divided into three groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5, and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3, while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2009]